

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

SO₂ absorption on K_xMn₈O₁₆ A, B, Cu-doped K_xMn₈O₁₆ B, and EMD
MnO₂ at 325°C




Fig. 1




Fig. 2a




Fig. 2b




Fig. 3a




Fig. 3b

SO₂ absorption on K_xMn₈O₁₆ B at 250°C, 325°C and 475°C
(~7500hr⁻¹ GHSV).




Fig. 4

Feed gas GHSV effect on the SO_2 absorption on $\text{K}_x\text{Mn}_8\text{O}_{16}$ B at 325°C




Fig. 5




Fig. 6




Fig. 7




Fig. 8




Fig. 9




Fig. 10




Fig. 11